

INFORMATION, EXPANSIVE NONDECELERATIVE UNIVERSE AND SUPERSTRING THEORY

Miroslav Súkeník and Jozef Šima

Slovak Technical University, Bratislava, Slovakia

Abstract. Stemming from relationships between a number of information describing a system and entropy content of the system it is possible to determine maximal cosmological time. The contribution manifests a compatibility of the superstring theory and the model of Expansive Nondecelerative Universe.

Entropy and Information

In spite of several justified objections raised to the second law of thermodynamics in specific cases [1], its validity for the Universe and the importance of entropy conception is generally accepted. Along with its probability meaning expressed by relation [2]

$$S = -k \cdot \ln p = k \cdot \ln P \quad (1)$$

where S is the entropy, k is the Boltzmann constant, p is the probability of the existence of a microstate, P is the number of microstates associated with a macroscopic state, entropy is also a measure of information needed to describe system properties. In accordance with a common usage, entropy will be expressed as dimensionless quantity, number of information is given in bit. Since a number of information transferred I by binary characters is [2, 3]

$$I = -\log_2 p \quad (2)$$

the entropy of a state fully describable by a number of information I is then

$$S = I(k \ln 2) \cong 9 \times 10^{-24} I \quad (3)$$

Thus, a number of information needed to fully describe a system can be expressed either via its entropy or by a number of bits.

Applying the above ideas to the Universe with a permanent increase of its entropy content, it is obvious that to describe such a universe, a number of information must grow as well.

Information, ENU and the Universe Dimensionality

One of the corner-stones of the holographic model of the Universe, elaborated by Bohm [4], declares that every point of the Universe (every elemental particle) is in mutual contact with the other points (particles) and holds the information about the whole Universe. The higher dimensionality of the Universe, the higher number of information is needed to its description. A postulate relating a maximum information $I_{\max}$ and a certain space dimensionality n of the Universe may be generally expressed as

$$\log_2 I_{\max} = 2^{(n-1)} \quad (4)$$

As a starting point (rationalized latter) let us suppose that

$$n = 10 \quad (5)$$

Then it follows from (4) and (5)

$$I_{\max} \cong 10^{154} \quad (6)$$

Corresponding maximum entropy based on (3) and (6) is

$$S_{\max} \cong 10^{131} \quad (7)$$

In the ENU model [5] it holds

$$a = c.t_c = \frac{2G.m_U}{c^2} \cong 1.3 \times 10^{26} \text{m} \quad (8)$$

where a is the gauge factor, t_c is the cosmological time, and m_U is the mass of the Universe. The elementary particles did not exist at the very beginning of the Universe, the gravitational field quanta, however, existed. This is why the entropy of the Universe can be expressed by means of a number of gravitational field quanta at a given cosmological time. If the mean energy of a gravitational field quantum is denoted as E_g , then

$$S = \frac{m_U.c^2}{|E_g|} = \frac{t_c.c^5}{2G.|E_g|} \quad (9)$$

As shown in our previous paper [6], it holds in the ENU

$$i.\hbar \frac{d\Psi_g}{dt} = E_g.\Psi_g \quad (10)$$

where Ψ_g is the wave function of the Universe defined as

$$\Psi_g = e^{-i(t_{Pc}.t_c)^{-1/2}.t} \quad (11)$$

It follows from (10) and (11) that

$$|E_g| = \hbar(t_{Pc}.t_c)^{-1/2} \quad (12)$$

where t_{Pc} is the Planck time

$$t_{Pc} = \left(\frac{G.\hbar}{c^5} \right)^{1/2} = 5.39056 \times 10^{-44} \text{s} \quad (13)$$

The entropy content at a time t is then, based on (9), (10), and (11), given by

$$S = \left(\frac{t}{t_{Pc}} \right)^{3/2} \quad (14)$$

i.e. at the time being

$$S \cong 10^{92} \quad (15)$$

A crucial conclusion stems from (7) and (14). It determines a cosmological time in which the maximum entropy should be reached

$$t_{c(\text{max})} \cong 10^{35} - 10^{36} \text{years} \quad (16)$$

A significance of this result lies in the fact that $t_{c(\text{max})}$ represents the time for which a decay of of baryonic matter is anticipated.

Concluding Remarks

Standard cosmological theories deals with the 3-dimensional space. Within the Kaluza-Klein theoretical approaches [7] more-dimensional spaces ($n = 4$ to 6) are elaborated [8]. A classic superstring theory used space with $n = 9$, in the M-theory $n = 10$. There are other conceptions within the superstring theories [9, 10] in which a number of dimensions is even higher.

The ENU model is compatible with superstring theory having $n = 10$. Calculations based on relations present in this work lead to a result showing that if $n < 10$, the maximum cosmological time would be less than the present time.

New results might be obtained when some features of the ENU model (such as the matter creation and gravitational energy localization) are incorporated into superstring theory.

References

1. V. Čápek, J. Bok, *Physica A* 290 (2001) 379
2. W. Yourgrau, A. van der Merwe, *Entropy, Information and Statistical Thermodynamics*, in *Cosmology, Fusion and Other Matters* (F. Reines, ed.), Adam Hilger, London, 1972, p.241
3. B.H. Mahan, *University Chemistry* (2nd ed.), Addison-Wesley Publ. Comp., London, 1972, p. 340
4. D. Bohm, B.J. Hiley, *The Undivided Universe*, Routledge, London, 1993
5. V. Skalský, M. Súkeník, *Astrophys. Space Sci.*, 178 (1991) 169
6. M. Súkeník, J. Šima, Preprint gr-qc/0010061
7. T. Appelquist, A. Chodos, P.G. Freund (Eds.) *Modern Kaluza-Klein Theories*, Addison-Wesley, Reading, MA., 1987
8. T. Gherghetta, M. Shaposhnikov, *Phys. Rev. Lett.*, 85 (2000) 240
9. M.B. Green, J.H. Schwarz, E. Witten, *Superstring Theory*, Vol. 1, 2. Cambridge University Press, Cambridge, 1989
10. M. Kaku, *Introduction to Superstrings*, Springer-Verlag, New York, 1988